

## **Quasi-Elastic Neutron Scattering**

Niina Jalarvo





#### How to become a neutron scattering scientist





#### **Overview**


#### INTRO to QENS

**QENS** Theory

**QENS** Instruments


Science Examples





# What is QENS

### Quasi Elastic Neutron Scattering QENS



- Quasi elastic neutron scattering is a limiting case of inelastic neutron scattering
- Doppler type of broadening of the elastic line due to a small energy transfer between the neutrons and the atoms in the sample



#### What can be measured


Probes slow dynamics in atomic length scale

- Translational diffusion
- Molecular reorientations
- Relaxation processes


Applicable to wide range of scientific topics

- Materials science: fuel cells, batteries, hydrogen storage,
- Soft Matter: polymer nanocomposites and blends, organic photovoltaics, polymer electrolytes
- Biology: hydration water, dynamics of proteins
- Chemistry: water interfaces, ionic liquids, clays, porous media, complex fluids, surface interactions


Results are comparable to Molecular nulations



Nature Communications 6, 7124 (2015)



Journal of Physical Chemistry C, 123, 2019 (2019).





Soft Matter, 7, 12, 5745-5755 (2011)



#### **Neutron-Material Interaction**

- Cross section (σ) Area related to the probability that a neutron will interact with a nucleus in a particular way (e.g. scattering or absorption)
- Light element sensitivity in presence of heavy elements
- Systems containing a reasonable proportion of H atoms, scattering from H tends to dominate
- Isotopic sensitivity
  - H-D contrast, H large incoherent cross-section
    - Use of deuteration/selective deuteration to suppress incoherent scattering
- Thermal neutron wavelengths (few Å's) are comparable to atomic / molecular distances
- Thermal neutron energies (few meV's) comparable to energies of excitation in materials
  - => vibrations, librations, reorientations, diffusion, and relaxational processes can be observed





# QENS theory


### **Scattering Kinematics**

The collision of two objects (e.g. neutron and sample atom) can be described in terms of momentum and energy conservation.

Neutron scattering events are described by means of energy and momentum transfer.

$$\begin{split} \hbar \vec{Q} &= \hbar \vec{k_1} - \hbar \vec{k_0} \\ \hbar \omega &= E_1 - E_0 \end{split}$$


$$\vec{p}_0, E_0$$
  $\vec{p}_1, E_1$   $2\theta$ 



$$\hbar\omega = 0$$
 — **ELASTIC** scattering

$$\hbar\omega \neq 0$$
 — INELASTIC scattering

$$\hbar\omega\approx 0$$
 — QUASIELASTIC scattering



#### Incoherent vs Coherent Neutron Scattering

Different atoms and isotopes have different coherent and incoherent scattering cross sections

Element	σ <sub>coh</sub> (barns)	σ <sub>inc</sub> (barns)
Hydrogen (H)	1.8	<mark>79.9</mark>
Deuterium (D)	5.6	2.0
Carbon (C)	5.6	0.001
Oxygen (O)	4.232	0

If the scattered neutron waves from the different nuclei have RANDOM relative phases (no interference)

=> INCOHERENT SCATTERING

If the scattered neutron waves from the different nuclei have definite relative phases, they can interfere

=> COHERENT SCATTERING

#### **DYNAMICS**

Protonated sample to observe single particle dynamics (quasielastic) and for the inelastic spectrum to weight hydrogen vibrations.

#### **STRUCTURE**

**Deuterated** sample to obtain structure and collective excitations.

Deuteration can help to suppress dynamics of particular groups



#### **QENS - Incoherent and Coherent Scattering**

 Large proportion of QENS experiments focus on dynamics of hydrogenous samples

$$S(Q, \varpi) = S_{inc}(Q, \varpi) + S_{coh}(Q, \varpi)$$

No information about structure Contaminates QENS signal (Bragg peaks)
Dominated by H dynamics

- Coherent QENS signal observable e.g. oxide ion diffusion
- Mixed coherent and incoherent QENS signal e.g. lithium or sodium diffusion



#### Scattering Function S(Q,ω)

The quantity measured in a neutron scattering experiment is the double differential cross-section

$$\frac{d^2\sigma}{d\Omega d\omega} \propto \frac{k_f}{k_i} \frac{\sigma}{4\pi} S(\boldsymbol{Q},\omega)$$

that gives the proportion of neutrons with an incident energy  $E_{\theta}$  scattered into a solid angle element  $d\Omega$  with an energy between  $E_{I}$  and  $E_{I}+dE_{I}$ .

#### Double differential cross-section:

$$\frac{\partial^2 \sigma}{\partial \Omega \delta E} = \left(\frac{\partial^2 \sigma}{\partial \Omega \delta E}\right)_{incoh} + \left(\frac{\partial^2 \sigma}{\partial \Omega \delta E}\right)_{coh}$$

 $\sigma$  = cross-section  $\Omega$  = solid angle, and E = energy transfer( $E = \hbar \omega$ )


#### COHERENT SCATTERING:


- Collective excitation from periodic samples
- Structure

#### INCOHERENT SCATTERING:

Vibrational modes from inelastic spectra


$$\frac{\partial^2 \sigma}{\partial \Omega \delta E} = \left(\frac{\partial^2 \sigma}{\partial \Omega \delta E}\right)_{incoh} = \frac{1}{4\pi N} \frac{\overrightarrow{k_f}}{\overrightarrow{k_i}} [\sigma_{incoh} S_{incoh}(Q, E)]$$







#### **QENS** spectra



Dynamic scattering function provides information on the sample states

Elastic intensity Debye-Waller factor: Vibrational amplitudes

Quasielastic intensity  $\longrightarrow$  A<sub>0</sub> = EISF (ratio elastic/total): Geometry of motion

Quasielastic broadening ------> Width: Characteristic time scale / diffusion



#### **QENS** Data Analysis

$$I(Q,E) = \underbrace{\left[p_1(Q)\delta(E) + \left(1 - p_1(Q)\right)S_{incoh}(Q,E)\right]}_{\text{Elastic}} \otimes R(Q,E) + \underbrace{B(Q,E)}_{\text{Resolution}}$$


$$S_{incoh}(Q,E) \equiv S_{QE}(Q,E)$$

$$S_{QE}(Q,E) = \frac{1}{\pi} \underbrace{\frac{DQ^2}{E^2 + (DQ^2)^2}}_{\text{FT-1 in time}} \text{In time}$$

$$I_s(Q,t) = \exp(-DQ^2t)$$

$$I_s(Q,t) = \exp(-DQ^2t)$$

$$G_s(r,t) = (4\pi Dt)^{-3/2} \exp(-r^2/4Dt)$$





Intermediate scattering function,  $I_{inc}(Q,t)$  can be obtained from a molecular dynamic simulation


#### **Elastic Window Scan**

- Elastic intensity scan as a function of temperature is a typical approach to estimate dynamic transitions.
- Resembles a DSC scan, i.e. locate transition temperature at which the dynamics enter the time window of the neutron spectrometer.
- Derive MSD using Gaussian approximation


$$S_{el}(Q,\omega) = A * e^{-Q^2 \langle r^2 \rangle / 3}$$








#### **QENS** diffusion models






- (TD) Translational Diffusion following Fick's law
- (CE) Chudley-Elliot -model, jump diffusion on a lattice
- (SS) Singwi-Sjölander -model, alternation between oscillatory motion and directed motion
- (HR) Hall-Ross –model, jump diffusion within a restricted volume

#### Spatially restricted diffusion

- Jumps between 2, 3, ... n sites
- Rotational diffusion on a circle
- Diffusion on a sphere
- Diffusion inside a sphere, cylinder

Angular dependency gives access to fundamental processes



### QENS Instruments

#### **QENS** Instruments

Currently about 20 QENS spectrometers in the world (in the U.S.,Germany, France, Switzerland, U.K. Japan and Australia)

### **Backscattering Spectrometers**

- High energy resolution
  - Resolution determined by the instrument (final energy of neutrons fixed)
  - Access to slower dynamics on nanosecond to picosecond time scale
  - Dynamic range limited


### Time-of-Fligth Spectrometers

- Lower energy resolution
  - Resolution can be varied by changing the energy of incoming neutrons
  - Access to dynamics on picosecond time scale
  - Larger dynamic range accessible



#### **QENS** and Neutron Scattering Instruments

- Dynamics in sample measurable
  - length scales set by Q range
    - $Q = 2\pi/d$
    - $0.1 \text{ Å}^{-1} < Q < 4 \text{ Å}^{-1} \rightarrow 60 \text{ Å} > d > 1.6 \text{ Å}$
  - Time scales set by the elastic energy resolution
    - higher resolution → longer times/slower motion (ns time scales accessible)
    - lower resolution → shorter times/faster motion (ps time scales accessible)
- interchange
  - dynamic range / resolution / count rate
  - Neutron λ vs Q
    - large λ -> high resolution -> long times/slow motions
    - large λ -> limited Q-range, limited length scales




M. Karlsson, Phys. Chem. Chem. Phys., 2015, 17, 26.





#### **BASIS** overview

BAckscattering SIlicon Spectrometer is a high-energy resolution, wide-dynamic range inverted geometry neutron spectrometer built on BL2 and facing a decoupled supercritical hydrogen, centerline-poisoned moderator




- Incident Flight Path 84 m moderatorsample position
  - Curved Guide: 10 cm wide x 12 cm tall, 1000 m radius of curvature, lineof-sight at 31 m
  - Straight Guide: 10 cm wide x 12 cm tall
  - Converging Funnel: last 7.7 m; exit 3.25 cm x 3.25 cm, stops 27.5 cm from sample
- Chopper System
  - 3 bandwidth/frame overlap choppers at 7, 9.25 and 50 m
  - Operation at 60 (standard), 30, 20, 15, 12, or 10 Hz
  - Bandwidth (full choppers transmission) of about 0.5 Å at 60 Hz

#### **Instrument Specifications**








	Si111	Si311
Elastic energy	2.08 meV	7.63 meV
Bandwidth 60 Hz 30 Hz	±100 μeV ±200 μeV	±660 μeV ±1700 μeV
Elastic resolution (HWHM)	3.6 μeV	15 <i>μ</i> eV
Q range (elastic)	$0.2  \text{Å}^{-1} < Q < 2.0  \text{Å}^{-1}$	$0.4 \text{ Å}^{-1} < Q < 3.8 \text{ Å}^{-1}$


- Radial Collimator restricts analyzer view of the sample
- Final Evacuated Flight Path - 2.5 m sample analyzer, ~ 2.23 m analyzer – detector
- Detector Choice LPSD <sup>3</sup>He tubes






**Instrument Description** 

- direct-geometry, multi-chopper inelastic / QENS spectrometer designed to provide flexibility in the choice of energy resolution
  - best at low incident energies (2 to 50 meV).
  - typical experiments use energy resolutions between 10 and 500  $\mu$ eV.





#### **Specifications**



Source-sample distance Sample-detector distance Angular coverage

Energy resolution
Incident energy range
Momentum transfer range

36.2 m

3.5 m

Horizontally: -50° - +140°


Vertically:±16°

 $10 - 500 \,\mu\text{eV}$ 

0.5 - 80 meV

0.05 - 10 Å - 1








# QENS - Science examples


#### Science Example 1: Molecular Reorientations


- Polyoligosilsesquioxane (POSS) Ligand Dynamics
- How do ligand dynamics contribute to the functionalities











#### Barrier for methyl rotation

$$V(\theta) = \frac{V_3}{2}(1 - \cos 3\theta)$$


$$H\psi_n(\theta) = E_n\psi_n(\theta)$$


 $-> V_3 = 74.4 \text{ meV } (7.18 \text{ kJ/mol})$ 



#### Science Example 2: Superionic Conductors

- Inorganic Na-ion superionic conductors play a vital role in all-solid-state Na batteries that operate at RT
- fundamental understanding of the Na-ion diffusion mechanisms in Na<sub>3</sub>SbS<sub>4</sub> with different crystal structures (e.g., tetragonal and cubic) from QENS
- The high degree of symmetry in cubic Na<sub>3</sub>SbS₄ leads to less interatomic correlations between Na and S(Sb) atoms, a shorter jump distance (2.85 Å), and a larger diffusion coefficient.





0

0.3

0.4 0.5

0.6


 $Q(\hat{A}^{-1})$ 

0.7



0.8 0.9 1.0


#### Science Example 3: Metal hydride dynamics



#### Softening of modes


H(1) and H(2) behave similarly with temperature


- •Modes merge with background at 600 K
  - Hydrogen releases and begins diffusing
  - ·Same temperature as onset of observable QENS diffusion






Science example 4: Elucidate Li<sup>+</sup> vs H<sup>+</sup> ion diffusion






H<sup>+</sup> ions were found to be immobile while Li<sup>+</sup> ion maintained a desired mobility in the solid electrolyte (Li<sub>6.25-x</sub>H<sub>x</sub>Al<sub>0.25</sub>)La<sub>3</sub>Zr<sub>2</sub>O<sub>12</sub>



 $Q^{2}(A^{-2})$ 



- □ Li+ ions show good ion mobility in the structure, the H+ ions are immobile at RT.
- Inactivity of H+ ions should contribute to the interfacial stability of LLZO being used as a protection layer for Limetal in aqueous Li batteries.
- ☐ This work provides a new method to probe diffusion behavior of different ions in solids that contain multiple mobile ion species





Science Example 5: Water Dynamics in Anion


**Exchange Membranes** 


Ionic conductivity and water transport are key properties for applications

- Depends on water content
- Water transport in AEM multiscale problem
  - Investigation of transport in multiple time- and length scales for structure – function insight













### Science example 6: $BaTiO_{3-x}H_x$ , Dynamics of Hydride Ions and The Role of Oxygen Vacancies

- hydride ion conducting material of interest for several electrochemical applications
- fundamental questions surrounding the defect chemistry and hydride ion transport mechanism



QENS jump diffusion of H **low T**  $\leq$ 250 K, jump distance ~2.8 Å, D ~ 0.4 × 10<sup>-6</sup> cm<sup>-2</sup> s<sup>-1</sup> at 225 K **high T** >400 K, jump distance ~ 4 Å D ~ 100 × 10<sup>-6</sup> cm<sup>-2</sup> s<sup>-1</sup> at 700 K  $E_{\rm s}$  = 0.1 eV



INS and DFT data confirm the presence of both nearest and next-nearest neighbouring oxygen vacancies to the hydride ions.

This supports QENS results, hydride ion transport is governed by jump diffusion dynamics between neighbouring and next-nearest neighbouring hydride ion-oxygen vacancy local structures.



#### **Summary:**

- QENS is a unique technique to measure diffusive dynamics providing exclusive information about the geometry of the diffusion
  - accessible through Q-dependence
  - Large range of time scales available depending of the selected QENS instrument (sub-picosecond < t < nanosecond (μsec for NSE)</li>
  - Hydrogen sensitivity
- Instrument selection is a critical decision
  - resolution to match the time scale of the diffusion process
  - Q range to match the diffusion length scale
- Suitable technique to study dynamics in large variety of materials and science problems.







Questions?

#### Literature:

- Quasielastic Neutron Scattering, M.
   Bee (Bristol, Adam Hilger, 1988).
- Quasielastic Neutron Scattering and Solid State Diffusion, R. Hempelmann (2000).

